

# An Estimate for Lipschitz Constants of Metric Projections\*

Chong Li

*Department of Applied Mathematics, Southeast University, Nanjing 210096, P. R. China*

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

*Department of Mathematics, Hangzhou University, Hangzhou 310028, P. R. China*

and

Wenshan Yang

*Department of Mathematics, Zhejiang Normal University, Jinhua 321004, P. R. China*

*Submitted by Joseph A. Ball*

Received March 2, 1998

The main purpose of the paper is to give a global estimate for Lipschitz constants of metric projections in a  $p$ -uniformly convex and  $q$ -uniformly smooth Banach space. © 1999 Academic Press

## 1. INTRODUCTION

Metric projection operators  $P_\Omega$  on convex closed sets  $\Omega$  (in the sense of best approximations) are widely used in theoretical and applied areas of mathematics, especially connected with problems of optimization and approximation. As examples one can consider iterative-projective methods for solving equations, variational inequalities and minimizations of functionals [1], and methods of alternating projections for finding common points of convex closed sets in Hilbert space [4, 5, 7].

\* Supported by National Natural Science Foundation of China.

It is well known [9] that in a Hilbert space  $X$  the following estimate holds:

$$\|P_{\Omega}(x) - P_{\Omega}(y)\| \leq \|x - y\|, \quad \text{for all } x, y \in X, \quad (1)$$

and many problems can be solved by applying the inequality (1). Therefore, one natural method to solve problems in Banach spaces  $X$  is to establish estimates analogous to (1). This is in fact shown by some author's recent work; see, for example, Xu and Roach [11], Alber and Notik [3], and Alber [2].

However, applying their results to a Hilbert space  $X$ , we can only obtain the following estimate:

$$\|P_{\Omega}(x) - P_{\Omega}(y)\| \leq c\|x - y\|, \quad \text{for all } x, y \in X, \quad (2)$$

for some constant  $c > 1$ . Comparing (2) with (1), it is natural to ask whether there exists a common estimate for  $\|P_{\Omega}(x) - P_{\Omega}(y)\|$  in a uniformly convex and uniformly smooth Banach space such that (1) holds for a Hilbert space. The purpose of the present paper is to give such an estimate for  $\|P_{\Omega}(x) - P_{\Omega}(y)\|$  in a  $p$ -uniformly convex and  $q$ -uniformly smooth Banach space.

## 2. PRELIMINARIES

Now let us recall the definition of the metric projection operator. Let  $\Omega$  be a convex closed subset in a Banach space  $X$ . For any  $x \in X$  an element  $\bar{x} \in \Omega$  such that  $\|x - \bar{x}\| = \min_{m \in \Omega} \|x - m\|$  is called a nearest point to  $x$ . If for each  $x \in X$ , it has a unique nearest point in  $\Omega$ , then define  $P_{\Omega}(x)$  to be the nearest point of  $x$ . It is known that  $P_{\Omega}$  is well defined in a uniformly convex Banach space. The moduli of convexity and smoothness of  $X$  are defined, respectively, by

$$\delta_X(\epsilon) = \inf\left\{1 - \left\|\frac{1}{2}(x + y)\right\| : \|x\| = \|y\|, \|x - y\| = \epsilon\right\}, \quad 0 \leq \epsilon \leq 2,$$

and

$$\rho_X(t) = \sup\left\{\frac{1}{2}(\|x + y\| + \|x - y\|) - 1 : \|x\| = 1, \|y\| = t\right\}, \quad t > 0.$$

Let  $p \geq 2$ ,  $1 < q \leq 2$ ,  $X$  is said to be  $p$ -uniformly convex (resp.  $q$ -uniformly smooth) if there is a constant  $d > 0$  such that  $\delta_X(\epsilon) \geq d\epsilon^p$  (resp.  $\rho_X(t) \leq dt^q$ ).

Let

$$d_p = \inf \left\{ \frac{(1/2)\|x\|^p + (1/2)\|y\|^p - \|(x+y)/2\|^p}{\|(x-y)/2\|^p} : \|x-y\| > 0 \right\},$$

$$c_q = \sup \left\{ \frac{(1/2)\|x\|^q + (1/2)\|y\|^q - \|(x+y)/2\|^q}{\|(x-y)/2\|^q} : \|x-y\| > 0 \right\}.$$

From [10] we have

**PROPOSITION 1.** *Let  $p > 1$ . Then a Banach space  $S$  is  $p$ -uniformly convex if and only if  $d_p > 0$ .*

**PROPOSITION 2.** *Let  $q > 1$ . Then a Banach space  $X$  is  $q$ -uniformly smooth if and only if  $c_q > 0$ .*

The following proposition is simple and well known.

**PROPOSITION 3.** *Let  $q > 1$  and  $X$  be strictly convex. Then for any  $x, y \in X$ ,  $f_x \in J_q(x)$ ,  $f_y \in J_q(y)$ , there holds*

$$\langle x - y, f_x - f_y \rangle > 0$$

where  $J_q(x) = \{x^* \in X^* : \langle x, x^* \rangle = \|x\|^q, \|x^*\| = \|x\|^{q-1}\}$ .

In order to give the main theorem we need a lemma about the constants  $d_p$  and  $c_q$ .

**LEMMA 1.** *Let  $X$  be a Banach space and  $n$  be a positive integer. Then for any  $x, y \in X$  the following hold*

$$\left\| \left(1 - \frac{1}{2^n}\right)x + \frac{1}{2^n}y \right\|^p \leq \left(1 - \frac{1}{2^n}\right)\|x\|^p + \frac{1}{2^n}\|y\|^p - d_p \left\{ \sum_{i=0}^{n-1} \left(\frac{1}{2^p}\right)^i \left(\frac{1}{2}\right)^{n-1-i} \right\} \left\| \frac{x-y}{2} \right\|^p \quad (3)$$

$$\left\| \left(1 - \frac{1}{2^n}\right)x + \frac{1}{2^n}y \right\|^q \geq \left(1 - \frac{1}{2^n}\right)\|x\|^q + \frac{1}{2^n}\|y\|^q - c_q \left\{ \sum_{i=0}^{n-1} \left(\frac{1}{2^q}\right)^i \left(\frac{1}{2}\right)^{n-1-i} \right\} \left\| \frac{x-y}{2} \right\|^q. \quad (4)$$

*Proof.* We will prove formula (3) by induction while formula (4) can be proved similarly. From the definition of  $d_p$ , we have that for any  $x, y \in X$ ,

$$\left\| \frac{1}{2}x + \frac{1}{2}y \right\|^p \leq \frac{1}{2}\|x\|^p + \frac{1}{2}\|y\|^p - d_p \left\| \frac{x-y}{2} \right\|^p.$$

This means that (3) holds for  $n = 1$ . Now assume that (3) holds for  $n = k$ . Let us prove that (3) holds for  $n = k + 1$ . In fact, for any  $x, y \in X$ ,

$$\begin{aligned} & \left\| \left(1 - \frac{1}{2^{k+1}}\right)x + \frac{1}{2^{k+1}}y \right\|^p \\ &= \left\| \frac{1}{2}x + \frac{1}{2} \left[ \left(1 - \frac{1}{2^k}\right)x + \frac{1}{2^k}y \right] \right\|^p \\ &\leq \frac{1}{2}\|x\|^p + \frac{1}{2} \left\| \left(1 - \frac{1}{2^k}\right)x + \frac{1}{2^k}y \right\|^p - d_p \left( \frac{1}{2^k} \right)^p \left\| \frac{x-y}{2} \right\|^p \\ &\leq \frac{1}{2}\|x\|^p + \frac{1}{2} \left[ \left(1 - \frac{1}{2^k}\right)\|x\|^p + \frac{1}{2^k}\|y\|^p \right] \\ &\quad - \frac{1}{2}d_p \left\{ \sum_{i=0}^{k-1} \left( \frac{1}{2^p} \right)^i \left( \frac{1}{2} \right)^{k-1-i} \right\} \left\| \frac{x-y}{2} \right\|^p - d_p \left( \frac{1}{2^p} \right)^k \left\| \frac{x-y}{2} \right\|^p \\ &= \left(1 - \frac{1}{2^{k+1}}\right)\|x\|^p + \frac{1}{2^{k+1}}\|y\|^p - d_p \left\{ \sum_{i=0}^k \left( \frac{1}{2^p} \right)^i \left( \frac{1}{2} \right)^{k-i} \right\} \left\| \frac{x-y}{2} \right\|^p. \end{aligned}$$

Hence (3) holds for  $n = k + 1$  and the proof is complete. ■

**PROPOSITION 4.** *Let  $q > 1$  and  $X$  be  $q$ -uniformly smooth. Then for any  $x, y \in X$  the following inequality holds*

$$\|x + y\|^q \leq \|x\|^q + q\langle y, J_q(x) \rangle + \frac{c_q}{2^{q-1} - 1} \|y\|^q. \quad (5)$$

*Proof.* From (4), it follows that for any  $n > 1$

$$\begin{aligned} & \frac{\|x + (1/2^n)y\|^q - \|x\|^q}{1/2^n} \\ &= \frac{\|(1 - (1/2^n))x + (1/2^n)(x + y)\|^q - \|x\|^q}{1/2^n} \\ &\geq \|x + y\|^q - \|x\|^q - \frac{c_q(1 - 2^{n-nq})}{2^{q-1} - 1} \|y\|^q. \end{aligned}$$

Thus we have that

$$\begin{aligned} q\langle y, J_q(x) \rangle &= \lim_{n \rightarrow \infty} \frac{\|x + (1/2^n)y\|^q - \|x\|^q}{1/2^n} \\ &\geq \|x + y\|^q - \|x\|^q - \frac{c_q}{2^{q-1} - 1} \|y\|^q. \end{aligned}$$

This proves the proposition. ■

### 3. MAIN THEOREM

Now we are ready to prove the main theorem

**THEOREM 1.** *Let  $X$  be a  $p$ -uniformly convex and  $q$ -uniformly smooth Banach space. Let  $\Omega$  be a closed convex subset in  $X$ . Then for any  $x, y \in X$  there holds*

$$\|P_\Omega x - P_\Omega y\| \leq \left[ \frac{pc_q(2^{p-1} - 1)}{qd_p(2^{q-1} - 1)} c^{p-q} \right] \|x - y\|^{q/p}$$

where  $c = \max\{\|x - P_\Omega y\|, \|y - P_\Omega x\|\}$ .

*Proof.* Let

$$Px = P_\Omega x, \quad Py = P_\Omega y,$$

and

$$x_n = Px + 2^n(x - Px)$$

Then  $Px_n = Px$  for any positive integer  $n$ . It follows that

$$\|x_n - Px\|^p \leq \|x_n - Py\|^p \leq \|x - Py + (2^n - 1)(x - Px)\|^p.$$

Dividing the inequality by  $2^n$  and using Lemma 1, we have

$$\frac{1}{2^n} \|x - Px\|^p \leq \frac{1}{2^n} \|x - Py\|^p - d_p \left\{ \sum_{i=0}^{n-1} \left( \frac{1}{2^p} \right)^i \left( \frac{1}{2} \right)^{n-1-i} \right\} \left\| \frac{Px - Py}{2} \right\|^p.$$

Multiplying the inequality by  $2^n$  and letting  $n \rightarrow \infty$ , we obtain

$$\frac{d_p}{2^{p-1} - 1} \|Px - Py\|^p \leq \|x - Py\|^p - \|x - Px\|^p.$$

Using Cauchy mean-valued theorem we obtain

$$\frac{d_p}{2^{p-1}-1} \|Px - Py\|^p \leq \frac{p}{q} c^{p-q} (\|x - Py\|^q - \|x - Px\|^q).$$

Similarly, we also have

$$\frac{d_p}{2^{p-1}-1} \|Px - Py\|^p \leq \frac{p}{q} c^{p-q} (\|y - Px\|^q - \|y - Py\|^q).$$

It follows that

$$2 \frac{d_p}{2^{p-1}-1} \|Px - Py\|^p \leq \frac{p}{q} c^{p-q} (\|x - Py\|^q - \|x - Px\|^q + \|y - Px\|^q - \|y - Py\|^q).$$

Note that (5) implies

$$\|x - Py\|^q - \|y - Py\|^q \leq q \langle x - y, J_q(y - Py) \rangle + \frac{c_q}{2^{q-1}-1} \|x - y\|^q,$$

$$\|y - Px\|^q - \|x - Px\|^q \leq q \langle y - x, J_q(x - Px) \rangle + \frac{c_q}{2^{q-1}-1} \|x - y\|^q.$$

Hence

$$\begin{aligned} & 2 \frac{d_p}{2^{p-1}-1} \|Px - Py\|^p \\ & \leq \frac{p}{q} c^{p-q} (q \langle x - y, J_q(y - Py) \rangle + q \langle y - x, J_q(x - Px) \rangle) \\ & \quad + 2 \frac{c_q}{2^{q-1}-1} \|x - y\|^q. \end{aligned}$$

From the characterization of a best approximation it follows that

$$\langle Px - Py, J_q(y - Py) \rangle \leq 0, \quad \langle Py - Px, J_q(x - Px) \rangle \leq 0.$$

This, together with Proposition 3, gives

$$\begin{aligned} & 2 \frac{d_p}{2^{p-1}-1} \|Px - Py\|^p \\ & \leq 2 \frac{p}{q} c^{p-q} \left( q \langle Qx - Qy, J_q(Qy) - J_q(Qx) \rangle + \frac{c_q}{2^{q-1}-1} \|x - y\|^q \right) \\ & \leq 2 \frac{p}{q} c^{p-q} \frac{c_q}{2^{q-1}-1} \|x - y\|^q \end{aligned}$$

where  $Qx = x - Px$ . That is,

$$\|Px - Py\| \leq \left[ \frac{pc_q(2^{p-1} - 1)}{qd_p(2^{q-1} - 1)} c^{p-q} \right] \|x - y\|^{q/p}.$$

The proof is complete. ■

If  $X$  is a Hilbert space then  $p = q = 2$  and  $d_p = c_q = 1$ . Therefore we have

**COROLLARY 1.** *Let  $X$  be a Hilbert space and  $\Omega$  be a convex closed subset in  $X$ . Then the following holds*

$$\|P_\Omega(x) - P_\Omega(y)\| \leq \|x - y\|, \quad \text{for all } x, y \in X.$$

**COROLLARY 2.** *Let  $X$  be the  $L_r(\mu)$ -space and  $\Omega$  be a convex closed subset in  $X$ . Then the following holds*

$$\|P_\Omega(x) - P_\Omega(y)\| \leq \begin{cases} \frac{r(r-1)c^{r-2}(2^{r-1} - 1)}{2} \|x - y\|^{2/r}, & r \geq 2 \\ \frac{2c^{2-r}}{r(r-1)(2^{r-1} - 1)} \|x - y\|^{r/2}, & r \leq 2. \end{cases}$$

*Proof.* We divide two cases,  $r \geq 2$  and  $r \leq 2$ , to prove the corollary.

(i) If  $r \geq 2$ , then  $p = r, q = 2$ , that is,  $L_r(\mu)$  is  $r$ -uniformly convex and 2-uniformly smooth. It follows from [6] that  $d_p = d_r = 1, c_q = c_2 \leq p - 1$ . Thus the result follows from Theorem 1.

(ii) If  $r \leq 2$ , then  $p = 2, q = r$ , that is,  $L_r(\mu)$  is 2-uniformly convex and  $r$ -uniformly smooth. From [8] we have  $d_p = d_2 \geq r - 1$ . Now let us show that  $c_q = c_r = 1$ .

For  $-1 \leq t < 1$ , let

$$f(t) = \frac{1/2 + (1/2)|t|^r - ((1+t)/2)^r}{((1-t)/2)^r}.$$

Then the derivative of  $f(t)$  for  $t \neq 0$  is

$$f'(t) = \frac{r(|t|^{r-1}s(t) - 2((1+t)/2)^{r-1} + 1)}{4((1-t)/2)^{r+1}}$$

where  $s(t) = t/|t|$  for  $t \neq 0$  and  $s(0) = 0$ . Write

$$g(t) = |t|^{r-1}s(t) - 2\left(\frac{1+t}{2}\right)^{r-1} + 1.$$

Then the derivative of  $g(t)$  for  $t \neq 0$  is

$$g'(t) = (r-1)(|t|^{r-2} - \left(\frac{1+t}{2}\right)^{r-2}).$$

Thus

$$\begin{aligned} g'(t) &> 0 & \forall t \in (-1/3, 0) \cup (0, 1) \\ g'(t) &< 0 & \forall t \in (-1, -1/3). \end{aligned}$$

From  $g(1) = g(-1) = 0$ ,  $g(0) < 0$  it follows that  $g(t) \leq 0$  for all  $t \in [-1, 1]$ . This implies that  $f'(t) \leq 0$  for all  $t \in [-1, 0) \cup (0, 1)$  so that

$$\sup\{f(t) : -1 \leq t < 1\} = f(-1) = 1.$$

Thus we have

$$\sup\left\{\frac{(1/2)|x|^r + (1/2)|y|^r - |(x+y)/2|^r}{|(x-y)/2|^r} : x \neq y\right\} = 1.$$

This implies that  $c_q = c_r = 1$  and the proof is complete. ■

## REFERENCES

1. Ya. I. Alber, Metric and generalized projection operators in Banach spaces: properties and applications, in "Theory and Applications of Nonlinear Operators of Monotone and Accretive Type" (A. Kartsatos, Ed.), pp. 15–50, Dekker, New York, 1996. E-preprint [funct-an/9311001](#).
2. Ya. I. Alber, A bound for modulus of continuity for metric projections in a uniformly convex and uniformly smooth space, *J. Approx. Theory* **85** (1996), 237–249.
3. Ya. I. Alber and A. I. Notik, On some estimates for projection operator in Banach space, *Comm. Appl. Nonlinear Anal.* **2** (1995), 47–56.
4. D. Butnariu and Y. Censor, Strong convergence of almost simultaneous block-iterative projection methods in Hilbert spaces, *J. Comput. Appl. Math.* **53** (1994), 33–42.
5. D. Butnariu and S. D. Flam, Strong convergence of expected-projection methods in Hilbert spaces, *Numer. Funct. Anal. Optim.* **16** (1995), 601–637.
6. J. A. Clarkson, Uniformly convex spaces, *Trans. Amer. Math. Soc.* **40** (1936), 396–414.



7. F. Deutsch, The method of alternating orthogonal projections, in "Approximation Theory, Spline Functions and Applications" (S. P. Singh, Ed.), pp. 105–121, Kluwer, Dordrecht, 1992.
8. T. C. Lim, H. K. Xu, and Z. B. Xu, An inequality and its applications to fixed point theory and approximation theory, in "Progress in Approximation Theory" (N. R. Pinkus *et al.*, Eds.), Academic, New York, 1992.
9. R. R. Phelps, Convex sets and nearest points, *Proc. Amer. Math. Soc.* **8** (1957), 790–797.
10. H. K. Xu, Inequalities in Banach spaces with applications, *Nonlinear Anal. Theory, Methods Appl.* **16** (1991), 1127–1138.
11. Z. B. Xu and G. F. Roach, On the uniform continuity of metric projections in Banach spaces, *Approx. Theory Appl.* **8** (1992), 11–19.